

Recent Progress on Tau Lepton Physics

A. Pich^a

^aDepartament de Física Teòrica, IFIC, Univ. València-CSIC, Apt. 22085, E-46071 València, Spain

Some important aspects of hadronic τ decays are reviewed: the determination of α_s from the inclusive τ hadronic width, the measurement of $|V_{us}|$ through the Cabibbo-suppressed decays of the τ , and the theoretical description of the $\tau \rightarrow \nu_\tau K\pi$ spectrum. The present status of other relevant electroweak topics, such as charged-current universality tests or bounds on lepton-flavour violation, has been already summarized in ref. [1].

1. The inclusive hadronic width of the tau

The hadronic τ decays turn out to be a beautiful laboratory for studying strong interaction effects at low energies [2, 3]. The τ is the only known lepton massive enough to decay into hadrons. Its semileptonic decays are then ideally suited for studying the hadronic weak currents.

The inclusive character of the total τ hadronic width renders possible an accurate calculation of the ratio [4–8]

$$R_\tau \equiv \frac{\Gamma[\tau^- \rightarrow \nu_\tau \text{ hadrons}]}{\Gamma[\tau^- \rightarrow \nu_\tau e^- \bar{\nu}_e]} = R_{\tau,V} + R_{\tau,A} + R_{\tau,S}.$$

The theoretical analysis involves the two-point correlation functions for the vector $V_{ij}^\mu = \bar{\psi}_j \gamma^\mu \psi_i$ and axial-vector $A_{ij}^\mu = \bar{\psi}_j \gamma^\mu \gamma_5 \psi_i$ colour-singlet quark currents ($i, j = u, d, s$):

$$\Pi_{ij,\mathcal{J}}^{\mu\nu}(q) \equiv i \int d^4x e^{iqx} \langle 0 | T(\mathcal{J}_{ij}^\mu(x) \mathcal{J}_{ij}^\nu(0)^\dagger) | 0 \rangle, \quad (1)$$

which have the Lorentz decompositions

$$\begin{aligned} \Pi_{ij,\mathcal{J}}^{\mu\nu}(q) &= (-g^{\mu\nu} q^2 + q^\mu q^\nu) \Pi_{ij,\mathcal{J}}^{(1)}(q^2) \\ &\quad + q^\mu q^\nu \Pi_{ij,\mathcal{J}}^{(0)}(q^2), \end{aligned} \quad (2)$$

where the superscript ($J = 0, 1$) denotes the angular momentum in the hadronic rest frame.

The imaginary parts of $\Pi_{ij,\mathcal{J}}^{(J)}(q^2)$ are proportional to the spectral functions for hadrons with the corresponding quantum numbers. The semi-hadronic decay rate of the τ can be written as an integral of these spectral functions over the

invariant mass s of the final-state hadrons:

$$\begin{aligned} R_\tau &= 12\pi \int_0^{m_\tau^2} \frac{ds}{m_\tau^2} \left(1 - \frac{s}{m_\tau^2}\right)^2 \\ &\quad \times \left[\left(1 + 2\frac{s}{m_\tau^2}\right) \text{Im}\Pi^{(1)}(s) + \text{Im}\Pi^{(0)}(s) \right]. \end{aligned} \quad (3)$$

The appropriate combinations of correlators are

$$\begin{aligned} \Pi^{(J)}(s) &\equiv |V_{ud}|^2 \left(\Pi_{ud,V}^{(J)}(s) + \Pi_{ud,A}^{(J)}(s) \right) \\ &\quad + |V_{us}|^2 \left(\Pi_{us,V}^{(J)}(s) + \Pi_{us,A}^{(J)}(s) \right). \end{aligned} \quad (4)$$

The contributions coming from the first two terms correspond to $R_{\tau,V}$ and $R_{\tau,A}$ respectively, while $R_{\tau,S}$ contains the remaining Cabibbo-suppressed contributions.

The integrand in Eq. (3) cannot be calculated at present from QCD. Nevertheless the integral itself can be calculated systematically by exploiting the analytic properties of the correlators $\Pi^{(J)}(s)$. They are analytic functions of s except along the positive real s -axis, where their imaginary parts have discontinuities. R_τ can then be written as a contour integral in the complex s -plane running counter-clockwise around the circle $|s| = m_\tau^2$:

$$\begin{aligned} R_\tau &= 6\pi i \oint_{|s|=m_\tau^2} \frac{ds}{m_\tau^2} \left(1 - \frac{s}{m_\tau^2}\right)^2 \\ &\quad \times \left[\left(1 + 2\frac{s}{m_\tau^2}\right) \Pi^{(0+1)}(s) - 2\frac{s}{m_\tau^2} \Pi^{(0)}(s) \right]. \end{aligned} \quad (5)$$

This expression requires the correlators only for complex s of order m_τ^2 , which is significantly larger than the scale associated with non-perturbative effects. Using the Operator Product

Expansion (OPE), $\Pi^{(J)}(s) = \sum_D C_D^{(J)} / (-s)^{D/2}$, to evaluate the contour integral, R_τ can be expressed as an expansion in powers of $1/m_\tau^2$. The uncertainties associated with the use of the OPE near the time-like axis are heavily suppressed by the presence in (5) of a double zero at $s = m_\tau^2$.

The combination $R_{\tau,V+A}$ can be written as [6]

$$R_{\tau,V+A} = N_C |V_{ud}|^2 S_{EW} \{1 + \delta_P + \delta_{NP}\}, \quad (6)$$

where $N_C = 3$ is the number of quark colours and $S_{EW} = 1.0201 \pm 0.0003$ contains the electroweak radiative corrections [9–11]. The dominant correction ($\sim 20\%$) is the perturbative QCD contribution δ_P , which is already known to $O(\alpha_s^4)$ [6, 12] and includes a resummation of the most important higher-order effects [7, 13].

Non-perturbative contributions are suppressed by six powers of the τ mass [6] and, therefore, are very small. Their numerical size has been determined from the invariant-mass distribution of the final hadrons in τ decay, through the study of weighted integrals [14],

$$R_\tau^{kl} \equiv \int_0^{m_\tau^2} ds \left(1 - \frac{s}{m_\tau^2}\right)^k \left(\frac{s}{m_\tau^2}\right)^l \frac{dR_\tau}{ds}, \quad (7)$$

which can be calculated theoretically in the same way as R_τ . The predicted suppression [6] of the non-perturbative corrections has been confirmed by ALEPH [15], CLEO [16] and OPAL [17]. The most recent analysis [18] gives

$$\delta_{NP} = -0.0059 \pm 0.0014. \quad (8)$$

The QCD prediction for $R_{\tau,V+A}$ is then completely dominated by δ_P ; non-perturbative effects being smaller than the perturbative uncertainties from uncalculated higher-order corrections. The result turns out to be very sensitive to the value of $\alpha_s(m_\tau^2)$, allowing for an accurate determination of the fundamental QCD coupling [5, 6]. The experimental measurement $R_{\tau,V+A} = 3.479 \pm 0.011$ implies [18]

$$\alpha_s(m_\tau^2) = 0.344 \pm 0.005_{\text{exp}} \pm 0.007_{\text{th}}. \quad (9)$$

The strong coupling measured at the τ mass scale is significantly larger than the values obtained at higher energies. From the hadronic decays of the Z , one gets $\alpha_s(M_Z^2) = 0.1191 \pm 0.0027$




Figure 1. Measured values of α_s at different scales. The curves show the energy dependence predicted by QCD, using $\alpha_s(m_\tau^2)$ as input. The corresponding extrapolated $\alpha_s(M_Z^2)$ values are shown at the bottom, where the shaded band displays the τ decay result within errors [18].

[12, 18, 19], which differs from $\alpha_s(m_\tau^2)$ by more than 20σ . After evolution up to the scale M_Z [20], the strong coupling constant in (9) decreases to [18]

$$\alpha_s(M_Z^2) = 0.1212 \pm 0.0011, \quad (10)$$

in excellent agreement with the direct measurements at the Z peak and with a better accuracy. The comparison of these two determinations of α_s in two very different energy regimes, m_τ and M_Z , provides a beautiful test of the predicted running of the QCD coupling; i.e., a very significant experimental verification of *asymptotic freedom*.

2. Perturbative contribution to R_τ

The recent calculation of the $\mathcal{O}(\alpha_s^4)$ contribution to $\Pi^{(0+1)}(s)$ [12] has triggered a renewed theoretical interest on R_τ [12, 18, 21, 22]. The perturbative contribution δ_P is extracted from the Adler

function

$$-s \frac{d}{ds} \Pi^{(0+1)}(s) = \frac{1}{4\pi^2} \sum_{n=0} K_n \left(\frac{\alpha_s(s)}{\pi} \right)^n. \quad (11)$$

For three flavours, the known coefficients take the values: $K_0 = K_1 = 1$; $K_2 = 1.63982$; $K_3(\overline{MS}) = 6.37101$ and $K_4(\overline{MS}) = 49.07570$ [12].

The perturbative component of R_τ is given by

$$\delta_P = \sum_{n=1} K_n A^{(n)}(\alpha_s), \quad (12)$$

where the functions [7]

$$A^{(n)}(\alpha_s) = \frac{1}{2\pi i} \oint_{|s|=m_\tau^2} \frac{ds}{s} \left(\frac{\alpha_s(-s)}{\pi} \right)^n \times \left(1 - 2 \frac{s}{m_\tau^2} + 2 \frac{s^3}{m_\tau^6} - \frac{s^4}{m_\tau^8} \right) \quad (13)$$

are contour integrals in the complex plane, which only depend on $a_\tau \equiv \alpha_s(m_\tau^2)/\pi$. Using the exact solution (up to unknown $\beta_{n>4}$ contributions) for $\alpha_s(s)$ given by the renormalization-group β -function equation, they can be numerically computed with a very high accuracy [7]. One can easily check that the results are very stable under changes of the renormalization scale and rather insensitive to the truncation of the β function (putting $\beta_4 = 0$ has a negligible impact). Thus, the resulting theoretical uncertainty on δ_P is small.

However if, instead of adopting the known values for $A^{(n)}(\alpha_s)$, one expands $\alpha_s(-s)$ in powers of $\alpha_s(m_\tau)$ inside the the integrals (13), the large logarithmic running along the circle $s = m_\tau^2 \exp(i\phi)$ ($\phi \in [0, 2\pi]$) gives rise to a nearly divergent series of the form $\delta_P = \sum_{n=1} (K_n + g_n) a_\tau^n$, where the g_n coefficients depend on $K_{m<n}$ and on $\beta_{m<n}$:

$$\delta^{(0)} = a_\tau + 5.20 a_\tau^2 + 26.4 a_\tau^3 + 127 a_\tau^4 + \dots \quad (14)$$

The “running” g_n contributions are much larger than the original K_n coefficients containing the Adler function dynamics ($g_2 = 3.563$, $g_3 = 19.99$, $g_4 = 78.00$) [7]. These generates a sizeable renormalization scale dependence, which is much larger than the naively expected $\mathcal{O}(\alpha_s^5)$ effect. The radius of convergence of this expansion is actually quite small. A numerical analysis of the series [7]

shows that, at the three-loop level, an upper estimate for the convergence radius is $a_{\tau, \text{conv}} < 0.11$, which is very close to the physical value. Thus, the fixed-order expansion (14) should not be used for accurate predictions of R_τ . The result (9) has been correctly obtained using Eq. (12) with the exact values of the functions $A^{(n)}(\alpha_s)$. The slightly different results quoted in refs. [12, 21] originate in their use of the pathological fixed-order expansion (14).¹

3. $|V_{us}|$ determination from tau decays

The separate measurement of the $|\Delta S| = 0$ and $|\Delta S| = 1$ tau decay widths provides a very clean determination of V_{us} [23, 24]. To a first approximation the Cabibbo mixing can be directly obtained from experimental measurements, without any theoretical input. Neglecting the small SU(3)-breaking corrections from the $m_s - m_d$ quark-mass difference, one gets:

$$|V_{us}|^{\text{SU}(3)} = |V_{ud}| \left(\frac{R_{\tau,S}}{R_{\tau,V+A}} \right)^{1/2} = 0.210 \pm 0.003.$$

We have used $|V_{ud}| = 0.97418 \pm 0.00027$ [25], $R_\tau = 3.640 \pm 0.010$ and the value $R_{\tau,S} = 0.1617 \pm 0.0040$ [24], which results from the most recent BaBar [26] and Belle [27] measurements of Cabibbo-suppressed tau decays [28]. The new branching ratios measured by BaBar and Belle are all smaller than the previous world averages, which translates into a smaller value of $R_{\tau,S}$ and $|V_{us}|$. For comparison, the previous value $R_{\tau,S} = 0.1686 \pm 0.0047$ [18] resulted in $|V_{us}|^{\text{SU}(3)} = 0.215 \pm 0.003$.

This rather remarkable determination is only slightly shifted by the small SU(3)-breaking con-

¹ A better convergence of the fixed-order expansion (14) is enforced in Ref. [21] through an artificial cancellation of the K_n and g_n contributions at higher orders. Since R_τ does not get corrections from $D = 4$ terms in the OPE, this behaviour is trivially accomplished assuming that the perturbative series is dominated by an $n = 2$ IR renormalon. While this provides an interesting academic model of higher-order contributions, the resulting wild behaviour of the Adler series is totally ad-hoc and generates problems for weighted distributions of the form (7). The non-perturbative correction in (8) would no longer be valid within this model, making the low value of $\alpha_s(m_\tau)$ claimed in [21] unjustified.

tributions induced by the strange quark mass. These corrections can be estimated through a QCD analysis of the differences [23, 24, 29–36]

$$\delta R_\tau^{kl} \equiv \frac{R_{\tau,V+A}^{kl}}{|V_{ud}|^2} - \frac{R_{\tau,S}^{kl}}{|V_{us}|^2}. \quad (15)$$

The only non-zero contributions are proportional to the mass-squared difference $m_s^2 - m_d^2$ or to vacuum expectation values of SU(3)-breaking operators such as $\delta O_4 \equiv \langle 0 | m_s \bar{s}s - m_d \bar{d}d | 0 \rangle \approx (-1.4 \pm 0.4) \cdot 10^{-3} \text{ GeV}^4$ [23, 29]. The dimensions of these operators are compensated by corresponding powers of m_τ^2 , which implies a strong suppression of δR_τ^{kl} [29]:

$$\delta R_\tau^{kl} \approx 24 S_{\text{EW}} \left\{ \frac{m_s^2(m_\tau^2)}{m_\tau^2} (1 - \epsilon_d^2) \Delta_{kl}(\alpha_s) - 2\pi^2 \frac{\delta O_4}{m_\tau^4} Q_{kl}(\alpha_s) \right\}, \quad (16)$$

where $\epsilon_d \equiv m_d/m_s = 0.053 \pm 0.002$ [37]. The perturbative corrections $\Delta_{kl}(\alpha_s)$ and $Q_{kl}(\alpha_s)$ are known to $O(\alpha_s^3)$ and $O(\alpha_s^2)$, respectively [29, 36].

The $J = 0$ contribution to $\Delta_{00}(\alpha_s)$ shows a rather pathological behaviour, with clear signs of being a non-convergent perturbative series. Fortunately, the corresponding longitudinal contribution to $\delta R_\tau \equiv \delta R_\tau^{00}$ can be estimated phenomenologically with a much better accuracy, $\delta R_\tau|^\perp = 0.1544 \pm 0.0037$ [23, 38], because it is dominated by far by the well-known $\tau \rightarrow \nu_\tau \pi$ and $\tau \rightarrow \nu_\tau K$ contributions. To estimate the remaining transverse component, one needs an input value for the strange quark mass. Taking the range $m_s(m_\tau) = (100 \pm 10) \text{ MeV}$ [$m_s(2 \text{ GeV}) = (96 \pm 10) \text{ MeV}$], which includes the most recent determinations of m_s from QCD sum rules and lattice QCD [38], one gets finally $\delta R_{\tau,th} = 0.216 \pm 0.016$, which implies [24]

$$\begin{aligned} |V_{us}| &= \left(\frac{R_{\tau,S}}{\frac{R_{\tau,V+A}}{|V_{ud}|^2} - \delta R_{\tau,th}} \right)^{1/2} \\ &= 0.2165 \pm 0.0026_{\text{exp}} \pm 0.0005_{\text{th}}. \end{aligned} \quad (17)$$

A larger central value, $|V_{us}| = 0.2212 \pm 0.0031$, is obtained with the old world average for $R_{\tau,S}$.

Sizeable changes on the experimental determination of $R_{\tau,S}$ are to be expected from the full

analysis of the huge BaBar and Belle data samples. In particular, the high-multiplicity decay modes are not well known at present. Thus, the result (17) could easily fluctuate in the near future. However, it is important to realize that the final error of the V_{us} determination from τ decay is completely dominated by the experimental uncertainties. If $R_{\tau,S}$ is measured with a 1% precision, the resulting V_{us} uncertainty will get reduced to around 0.6%, i.e. ± 0.0013 , making τ decay the best source of information about V_{us} .

An accurate measurement of the invariant-mass distribution of the final hadrons could make possible a simultaneous determination of V_{us} and the strange quark mass, through a correlated analysis of several weighted differences δR_τ^{kl} . The extraction of m_s suffers from theoretical uncertainties related to the convergence of the perturbative series $\Delta_{kl}(\alpha_s)$, which makes necessary a better understanding of these corrections.

4. $\tau \rightarrow \nu_\tau K \pi$ and $K \rightarrow \pi l \bar{\nu}_l$

The decays $\tau \rightarrow \nu_\tau K \pi$ probe the same hadronic form factors investigated in K_{l3} processes, but they are sensitive to a much broader range of invariant masses. A theoretical understanding of the form factors can be achieved, using analyticity, unitarity and some general properties of QCD, such as chiral symmetry and the short-distance asymptotic behaviour [2, 3].

Figure 2 compares the resulting theoretical description of the τ decay spectrum [39] with the recent Belle measurement [27]. At low values of s there is clear evidence of the scalar contribution, which was predicted previously using a careful analysis of $K\pi$ scattering data [38, 40]. From the measured τ spectrum one obtains $M_{K^*} = 895.3 \pm 0.2 \text{ MeV}$ and $\Gamma_{K^*} = 47.5 \pm 0.4 \text{ MeV}$ [39]. Since the absolute normalization is fixed by K_{l3} data to be $|V_{us}| f_+^{K^0 \pi^-}(0) = 0.21664 \pm 0.00048$ [41], one gets then a theoretical prediction for the branching fraction, $\text{Br}(\tau^- \rightarrow \nu_\tau K_S \pi^-) = 0.427 \pm 0.024\%$, in good agreement with the Belle measurement $0.404 \pm 0.013\%$, although slightly larger.

The τ determination of the vector form factor $f_+^{K\pi}(s)$ [39, 42] provides precise values for its slope and curvature, $\lambda'_+ = (25.2 \pm 0.3) \cdot 10^{-3}$ and

$\lambda''_+ = (12.9 \pm 0.3) \cdot 10^{-4}$ [39], in agreement but more precise than the corresponding K_{l3} measurements [41].




Figure 2. Theoretical description [39] (solid line) of the Belle $\tau^- \rightarrow \nu_\tau K_S \pi^-$ data [27]. The K^{*} (dashed-dotted) and scalar (dotted) contributions are also shown.

Acknowledgements

This work has been supported by MICINN, Spain (grants FPA2007-60323 and Consolider-Ingenio 2010 CSD2007-00042, CPAN) by the EU Contract MRTN-CT-2006-035482 (FLAVIANet) and by Generalitat Valenciana (PROMETEO/2008/069).

REFERENCES

1. A. Pich, Nucl. Phys. B (Proc. Suppl.) 181-182 (2008) 300; 169 (2007) 393.
2. A. Pich, Int. J. Mod. Phys. A 21 (2006) 5652.
3. A. Pich, *Tau Physics*, in *Heavy Flavours II*, eds. A.J. Buras and M. Lindner, Advanced Series on Directions in High Energy Physics 15 (World Scientific, Singapore, 1998) p. 453, arXiv:hep-ph/9704453.
4. E. Braaten, Phys. Rev. Lett. 60 (1988) 1606; Phys. Rev. D 39 (1989) 1458.
5. S. Narison and A. Pich, Phys. Lett. B 211 (1988) 183.
6. E. Braaten, S. Narison and A. Pich, Nucl. Phys. B 373 (1992) 581.
7. F. Le Diberder and A. Pich, Phys. Lett. B 286 (1992) 147.
8. A. Pich, Nucl. Phys. B (Proc. Suppl.) 39B,C (1995) 326.
9. W.J. Marciano and A. Sirlin, Phys. Rev. Lett. 61 (1988) 1815.
10. E. Braaten and C.S. Li, Phys. Rev. D 42 (1990) 3888.
11. J. Erler, Rev. Mex. Phys. 50 (2004) 200.
12. P.A. Baikov, K.G. Chetyrkin and J.H. Kühn, Phys. Rev. Lett. 101 (2008) 012002.
13. A.A. Pivovarov, Z. Phys. C 53 (1992) 461.
14. F. Le Diberder and A. Pich, Phys. Lett. B 289 (1992) 165.
15. ALEPH Collaboration, Phys. Rep. 421 (2005) 191; Eur. Phys. J. C 4 (1998) 409; Phys. Lett. B 307 (1993) 209.
16. CLEO Collaboration, Phys. Lett. B 356 (1995) 580.
17. OPAL Collaboration, Eur. Phys. J. C 7 (1999) 571.
18. M. Davier et al., Rev. Mod. Phys. 78 (2006) 1043; Eur. Phys. J. C 56 (2008) 305.
19. The LEP Collaborations ALEPH, DELPHI, L3 and OPAL and the LEP Electroweak Working Group, arXiv:0712.0929 [hep-ex]; <http://www.cern.ch/LEPEWWG/>.
20. G. Rodrigo, A. Pich and A. Santamaria, Phys. Lett. B 424 (1998) 367.
21. M. Beneke and M. Jamin, JHEP 0809 (2008) 044.
22. K. Maltman and T. Yavin, arXiv:0807.0650 [hep-ph].
23. E. Gámiz et al., Phys. Rev. Lett. 94 (2005) 011803; JHEP 0301 (2003) 060.
24. E. Gámiz et al., PoS KAON 008 (2007).
25. C. Amsler et al., *The Review of Particle Physics*, Phys. Lett. B 667, 1 (2008).
26. BaBar Collaboration, Phys. Rev. Lett. 100 (2008) 011801; Phys. Rev. D 76 (2007) 051104.
27. Belle Collaboration, Phys. Lett. B 654 (2007) 65; 643 (2006) 5.
28. S. Banerjee, PoS KAON 009 (2007).
29. A. Pich and J. Prades, JHEP 9910 (1999) 004; 9806 (1998) 013.

- 30. S. Chen et al., Eur. Phys. J. C 22 (2001) 31.
M. Davier et al., Nucl. Phys. B (Proc. Suppl.) 98 (2001) 319.
- 31. K.G. Chetyrkin, J.H. Kühn and A.A. Pivovarov, Nucl. Phys. B 533 (1998) 473.
- 32. J.G. Körner, F. Krajewski and A.A. Pivovarov, Eur. Phys. J. C 20 (2001) 259.
- 33. K. Maltman and C.E. Wolfe, Phys. Lett. B 639 (2006) 283. K. Maltman et al. arXiv:0807.3195 [hep-ph].
- 34. J. Kambor and K. Maltman, Phys. Rev. D 62 (2000) 093023; 64 (2001) 093014.
- 35. K. Maltman, Phys. Rev. D 58 (1998) 093015.
- 36. P.A. Baikov, K.G. Chetyrkin and J.H. Kühn, Phys. Rev. Lett. 95 (2005) 012003.
- 37. H. Leutwyler, Phys. Lett. B 378 (1996) 313.
- 38. M. Jamin, J.A. Oller and A. Pich, Phys. Rev. D 74 (2006) 074009.
- 39. M. Jamin, A. Pich and J. Portolés, Phys. Lett. B 640 (2006) 176; 664 (2008) 78.
- 40. M. Jamin, J.A. Oller, and A. Pich, Nucl. Phys. B 587 (2000) 331; 622 (2002) 279; Eur. Phys. J. C 24 (2002) 237.
- 41. FlaviaNet Working Group on Kaon Decays, Nucl. Phys. B (Proc. Suppl.) 181–182 (2008) 83.
- 42. D.R. Boito et al. arXiv:0807.4883 [hep-ph].